

---

**PHYSICS**

**9702/23**

Paper 2 AS Level Structured Questions

**October/November 2017**

MARK SCHEME

Maximum Mark: 60

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is a registered trademark.

---

This document consists of **7** printed pages.

| Question | Answer  | Marks     |
|----------|---|-----------|
| 1(a)(i)  | work (done) / time (taken) <b>or</b> energy (transferred) / time (taken)  | <b>B1</b> |
| 1(a)(ii) | Correct substitution of base units of all quantities into any correct equation for power.<br><br>Examples:<br><br>$(P = E / t \text{ or } W / t \text{ gives}) \text{ kg m}^2 \text{ s}^{-2} / \text{s} = \text{kg m}^2 \text{ s}^{-3}$<br><br>$(P = Fs / t \text{ or } mgh / t \text{ gives}) \text{ kg m s}^{-2} \text{ m} / \text{s} = \text{kg m}^2 \text{ s}^{-3}$<br><br>$(P = \frac{1}{2}mv^2 / t \text{ gives}) \text{ kg (m s}^{-1})^2 / \text{s} = \text{kg m}^2 \text{ s}^{-3}$<br><br>$(P = Fv \text{ gives}) \text{ kg m s}^{-2} \text{ m s}^{-1} = \text{kg m}^2 \text{ s}^{-3}$<br><br>$(P = VI \text{ gives}) \text{ kg m}^2 \text{ s}^{-2} \text{ A}^{-1} \text{ s}^{-1} \text{ A} = \text{kg m}^2 \text{ s}^{-3}$ | <b>A1</b> |
| 1(b)(i)  | units of $A$ : $\text{m}^2$ <b>and</b> units of $T$ : $\text{K}$  | <b>C1</b> |
|          | units of $k$ : $\text{kg m}^2 \text{ s}^{-3} / \text{m}^2 \text{ K}^4$<br>$= \text{kg s}^{-3} \text{ K}^{-4}$   | <b>A1</b> |
| 1(b)(ii) | curve from the origin with increasing gradient  | <b>B1</b> |

| Question | Answer  | Marks     |
|----------|---|-----------|
| 2(a)     | $\rho = m / V$ <b>or</b> $\rho = m / Ah$                                    | <b>B1</b> |
|          | $p = F / A$ <b>or</b> $p = W / A$   | <b>B1</b> |
|          | $p = [\rho Ahg] / A$ <b>or</b> $p = [\rho Vg] / [V / h]$ (so) $p = \rho gh$ | <b>A1</b> |

| Question | Answer  | Marks |
|----------|---|-------|
| 2(b)(i)  | 1. weight/gravitational (force)<br>upthrust (force)/buoyancy (force)<br>drag/viscous/frictional (force)/fluid resistance/resistance   | B1    |
|          | 2. weight = upthrust + viscous (force)  | B1    |
| 2(b)(ii) | <ul style="list-style-type: none"> <li>decrease in (gravitational) potential energy (of sphere) due to decrease in height (since <math>E_p = mgh</math>)</li> <li>increase in thermal energy due to work done against viscous force/drag</li> <li>loss/change of (total) <math>E_p</math> equal to gain/change in thermal energy</li> </ul> Any 2 points. | B2    |
| 2(c)(i)  | atmospheric pressure = $9.1(0) \times 10^4$ Pa  | A1    |
| 2(c)(ii) | $(\Delta)p = \rho g(\Delta)h$<br>$(9.15 - 9.10) \times 10^4 = \rho \times 9.81 \times (0.17 - 0.10)$  | C1    |
|          | $\rho = 730$ (728) $\text{kg m}^{-3}$   | A1    |

| Question | Answer  | Marks |
|----------|---|-------|
| 3(a)     | <u>sum/total</u> momentum (of system of bodies) is constant<br>or<br><u>sum/total</u> momentum before = <u>sum/total</u> momentum after                       | M1    |
|          | for an isolated system/no (resultant) <u>external</u> force   | A1    |
| 3(b)(i)  | $p = mv$  | C1    |
|          | $(4.0 \times 6.0 \times \sin \theta) - (12 \times 3.5 \times \sin 30^\circ) = 0$<br>or<br>$(m_A v_A \times \sin \theta) - (m_B v_B \times \sin 30^\circ) = 0$ | M1    |
|          | $\theta = 61^\circ$   | A1    |

| Question  | Answer  | Marks     |
|-----------|---|-----------|
| 3(b)(ii)  | shows the horizontal <u>momentum</u> component of ball A or of ball B as $(4.0 \times 6.0 \times \cos \theta)$ <b>or</b> $(12 \times 3.5 \times \cos 30^\circ)$ | <b>C1</b> |
|           | $(4.0 \times 6.0 \times \cos 61^\circ) + (12 \times 3.5 \times \cos 30^\circ) = 4.0v$ so $v = 12 \text{ (ms}^{-1}\text{)}$                                      | <b>A1</b> |
| 3(b)(iii) | initial $E_K (= \frac{1}{2} \times 4.0 \times 12^2) = 290 \text{ (288) (J)}$  | <b>M1</b> |
|           | final $E_K (= \frac{1}{2} \times 4.0 \times 6.0^2 + \frac{1}{2} \times 12 \times 3.5^2) = 150 \text{ (145.5) (J)}$  | <b>M1</b> |
|           | (initial $E_K >$ final $E_K$ ) so inelastic [both M1 marks required to award this mark]   | <b>A1</b> |

| Question | Answer   | Marks     |
|----------|--|-----------|
| 4(a)     | displacement of particles/vibration(s)/oscillation(s) is parallel to/along the direction of energy/propagation | <b>B1</b> |
| 4(b)     | period = $1 / 800 (= 1.25 \times 10^{-3} \text{ s})$   | <b>C1</b> |
|          | time-base setting = $1.25 \times 10^{-3} / 2.5$  | <b>C1</b> |
|          | $= 5.0 \times 10^{-4} \text{ s cm}^{-1}$   | <b>A1</b> |
| 4(c)(i)  | $I \propto A^2$  | <b>C1</b> |
|          | $(I_X / I_Y =) [r_Y / r_X]^2 = [A_X / A_Y]^2$  | <b>C1</b> |
|          | ratio $A_Y / A_X = 120 / 30$<br>$= 4.0$  | <b>A1</b> |

| Question | Answer  | Marks |
|----------|---|-------|
| 4(c)(ii) | 1. $v = f\lambda$   | C1    |
|          | minimum $\lambda = 330 / (800 + 16) = 0.40 \text{ m}$             | A1    |
|          | 2. $f_o / f_s = v / (v - v_s)$<br>$816 / 800 = 330 / (330 - v_s)$ | C1    |
|          | $v_s = 6.5 \text{ m s}^{-1}$                                      | A1    |

| Question  | Answer   | Marks |
|-----------|--|-------|
| 5(a)      | force <u>per</u> unit positive charge  | B1    |
| 5(b)(i)   | $s = \frac{1}{2}at^2$  | C1    |
|           | $a = (2 \times 0.045) / (1.5 \times 10^{-7})^2 = 4(.0) \times 10^{12} \text{ m s}^{-2}$      | A1    |
| 5(b)(ii)  | $F = 1.67 \times 10^{-27} \times 4.0 \times 10^{12} = 6.7 (6.68) \times 10^{-15} \text{ N}$  | A1    |
| 5(b)(iii) | 1. $E = F / Q$   | C1    |
|           | $= 6.68 \times 10^{-15} / 1.6 \times 10^{-19}$<br>$= 4.2 (4.18) \times 10^4 \text{ NC}^{-1}$ | A1    |
|           | 2. $E = V / d$   | C1    |
|           | $V = 4.18 \times 10^4 \times 0.045$<br>$= 1.9 \times 10^3 \text{ V}$                         | A1    |

| Question | Answer   | Marks |
|----------|--|-------|
| 5(c)     | $a = Eq / m$<br>or<br>$F = ma$ and $F = Eq$  | C1    |
|          | ratio = $\frac{(2 \times 1.6 \times 10^{-19}) \times (1.67 \times 10^{-27})}{(1.6 \times 10^{-19}) \times (4 \times 1.66 \times 10^{-27})}$ or $\frac{2 \times 1}{1 \times 4}$<br><br>= 0.50 | A1    |

| Question | Answer  | Marks |
|----------|---|-------|
| 6(a)(i)  | $P = VI$  | C1    |
|          | $I = 30 / 120$<br><br>= 0.25 A  | A1    |
| 6(a)(ii) | $Q = 0.25 \times 3.0 \times 3600 (= 2700)$  | C1    |
|          | number = $(0.25 \times 3.0 \times 3600) / 1.60 \times 10^{-19}$<br><br>= $1.7 \times 10^{22}$ | A1    |
| 6(b)     | $R = V / I$ or $R = P / I^2$ or $R = V^2 / P$   | C1    |
|          | = $120 / 0.25$ or = $30 / 0.25^2$ or = $120^2 / 30$ = $480 \Omega$                            | A1    |

| Question | Answer  | Marks     |
|----------|---|-----------|
| 6(c)     | $R = \rho l / A$  | <b>C1</b> |
|          | $A = (6.1 \times 10^{-7} \times 580 \times 10^{-3}) / 480 (= 7.37 \times 10^{-10})$     | <b>C1</b> |
|          | $d = [(4 \times 7.37 \times 10^{-10}) / \pi]^{1/2}$<br>$= 3.1 \times 10^{-5} \text{ m}$ | <b>A1</b> |
| 6(d)     | temperature decreases and so resistance decreases                                       | <b>B1</b> |

| Question | Answer   | Marks     |
|----------|--|-----------|
| 7(a)     | nucleons = 23  | <b>B1</b> |
|          | neutrons = 11  | <b>B1</b> |
| 7(b)     | similarity:<br>same (rest) mass<br><b>or</b><br>equal (magnitude of) charge  | <b>B1</b> |
|          | difference:<br>opposite (sign of) charge<br><b>or</b><br>one is matter and one is antimatter<br><b>or</b><br>one is an electron and one is an antielectron | <b>B1</b> |