Please check the examination deta	ils bel	ow before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel	Cen	tre Number	Candidate Number
Level 1/Level 2 GCSE (9-1)			
100011/100011 0001(5-1)			
Time 1 hour 45 minutes		Paper reference	1PH0/1F
Physics			
PAPER 1			
			Foundation Tier
(Very ground become			Tarritte
You must have: Calculator, ruler			Total Marks
Carcalatol, Falci			

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- A list of equations is included at the end of this exam paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

(a)	The S	un is	s at the centre of our Solar System.	
	The p	lane	ts orbit the Sun.	
	Which	n pla	net is nearest to the Sun?	(4)
	×	Α	Jupiter	(1)
	×	В	Mars	
	×	C	Mercury	
	×	D	Venus	
(b)	The M	loor	orbits the Earth.	
	Which	of t	these describes the Moon?	(1)
	\times	Α	an asteroid	(1)
	×	В	a comet	
	X	C	a nebula	
	\times	D	a natural satellite	
(c)	Any o	bjec	t weighs less on the Moon than it does on the Earth.	
				7/
				(2)
		Which Which Which Any of The g field s Sugge	Which pla A B C D b) The Moon Which of the A B C D c) Any object The gravit field strend Suggest the A	B Mars C Mercury D Venus b) The Moon orbits the Earth. Which of these describes the Moon? A an asteroid B a comet C a nebula



(d) The gravitational field strength on the Moon is 1.6 N/kg. The mass of a rock on the Moon is 6.0 kg.

Calculate the weight of this rock on the Moon.

State the unit of weight.

Use the equation

 $weight = mass \times gravitational \ field \ strength$

(3)

weight of rock =unitunit

(Total for Question 1 = 7 marks)



2 (a) Figure 1 shows the parts of the electromagnetic spectrum.

gamma rays	x-rays	J	visible	K	micro- waves	L
---------------	--------	---	---------	---	-----------------	---

Figure 1

(i) Which row of the table names the parts **J**, **K** and **L** of the electromagnetic spectrum?

		J	K	L
\times	A	infrared	radio	ultraviolet
\times	В	radio	infrared	ultraviolet
\times	C	ultraviolet	infrared	radio
\boxtimes	D	ultraviolet	radio	infrared

(ii) All electromagnetic waves can travel in a vacuum.

Which of these is the same for all electromagnetic waves travelling in a vacuum?

■ A amplitude

■ B frequency

D wavelength

- (b) X-rays can be useful and harmful to humans.
 - (i) State **one** way that x-rays are useful to humans.

(1)

(1)

(ii) State **one** way that x-rays are harmful to humans.

(1)



(c) A person warms their hands in front of a hot fire as shown in Figure 2.



(Source: © Andreas Saldavs/Shutterstock)

Figure 2

Use words from the box to complete the following sentences.

chemical infrared radio thermal ultraviolet

(2)

The electromagnetic waves that the fire mostly emits are waves.

These waves transfer energy to the hands.

(Total for Question 2 = 6 marks)



3	This question is about nuclea	ar reactions.	
		235) nucleus absorbs a neutron, the nucleus splits into	smaller narts
	This reaction is called nuc		Silianei pares.
		lei are produced from the fission of one nucleus of	
	uranium-235 (U-235)?		(1)
	🖾 A two		(- /
	■ B three		
	D six		
	(b) (i) When a uranium-235	(U-235) nucleus splits, neutrons are also emitted.	
	The neutrons may sta	rt a chain reaction.	
	Describe what is mea	nt by a chain reaction.	
	You may draw a diagr	ram to help with your answer.	(2)
			(2)
	(ii) In the nuclear reactor	of a power station, the chain reaction has to be control	olled.
	Explain the action of	a moderator in a nuclear reactor.	(2)
			(2)



(iii) In a nuclear reactor there are 2.0	X	< 10 ¹⁷ fission reactions each second.
--	---	---

Each fission reaction releases $4.0 \times 10^{-11} \text{J}$ of energy.

Calculate the energy released in 1 second.

(2)

energy released in 1 second =

(c) Nuclear fusion is a nuclear reaction that takes place in the Sun.

Describe what happens during a nuclear fusion reaction.

(2)

(Total for Question 3 = 9 marks)





- **4** (a) A cyclist has a mass of 64 kg.
 - (i) The cyclist rides from a flat road to the top of a hill.

The top of the hill is 24 m above the flat road.

Calculate the gain in gravitational potential energy, Δ GPE, of the cyclist.

Use $g = 10 \,\text{N/kg}$

Use the equation

$$\Delta GPE = m \times g \times \Delta h$$

(2)

(ii) The cyclist returns to the flat road.

The mass of the cyclist is 64 kg.

Calculate the kinetic energy of the cyclist when the cyclist is travelling at 6.0 m/s.

Use the equation

$$KE = \frac{1}{2} \times m \times v^2$$

(3

kinetic energy =

(iii) The cyclist then uses the brakes on the bicycle to stop.

Explain what happens to the kinetic energy of the cyclist.

(2)

(b) A different cyclist uses a motorised bicycle.

The motorised bicycle is powered by an electric motor.

Figure 3 is an energy diagram for the motor.




Figure 3

(i) Calculate how much energy is wasted.

(1)

energy wasted =



(ii) Calculate the efficiency of the electric motor.

(2)

Use the equation:

efficiency =
$$\frac{\text{(useful energy transferred by the device)}}{\text{(total energy supplied to the device)}}$$

efficiency of electric motor =

(Total for Question 4 = 10 marks)

5 (a) A car is travelling at 10 m/s.

The driver sees a danger and stops the car.

(i) The stopping distance for the car would be smaller if the car

(1)

- A had more passengers
- **B** had worn tyres
- C needed new brakes
- **D** was travelling more slowly

Figure 4 shows a speed-time graph for the driver stopping the car.




Figure 4

(ii) Use the graph to find the driver's reaction time.

(2)

reaction time =s

(2)

(b) Figure 5 shows the apparatus a student uses to investigate how the stopping distance of a toy car depends on the type of surface that it is stopping on.




Figure 5

Describe an experiment to find out how the stopping distance depends on the surface that stops the toy car.



(c) Figure 6 shows a set of results used to find the average stopping distance of the toy car on a surface.

test number	stopping distance in m
1	0.35
2	0.32
3	0.52
4	0.38
5	0.33

Figure 6

(i) State the anomalous value of stopping distance given in the table in Figure 6.

(1)

(ii) Use the results in Figure 6 to calculate the average stopping distance.

(2)

average stopping distance = m

(iii) State **one** way the student could increase the speed of the car as it reaches the flat surface.

(1)

(d) A car is travelling down a slope at 2.0 m/s.

The car accelerates for 4.0 s.

The speed of the car increases to 12 m/s.

Calculate the acceleration of the car.

Use the equation

$$a = \frac{(v - u)}{t}$$

(2)

acceleration of the car =m/s

(Total for Question 5 = 11 marks)



- **6** Quantities can be either scalar or vector.
 - (a) Which of these is a vector quantity?

(1)

- A mass
- **B** force
- **D** distance
- (b) Figure 7 shows a ball bearing as it falls slowly through a clear, dense liquid.




Figure 7

The apparatus in Figure 7 is used to find the average speed of the ball bearing as it falls.

(i) Devise an experiment to determine the average speed of the ball bearing as it falls through the liquid.

(4)

You should include:

- any extra apparatus you would use to take measurements
- the measurements you would take
- how you would calculate the speed.

(ii) A student thinks that the ball bearing falls through the liquid at a constant speed.

Explain how you could develop this experiment to determine if the ball bearing falls through the liquid at constant speed.

You may draw a diagram to help your answer.

(2)

(c) The ball bearing is now dropped through air.

The initial velocity of the ball bearing is zero.

The acceleration of the ball bearing is 10 m/s².

The ball bearing falls 1.5 m.

Calculate the velocity of the ball bearing when it has fallen 1.5 m.

Use the equation

$$v^2 - u^2 = 2 \times a \times x$$

(2)

velocity of ball bearing =m/s

(Total for Question 6 = 9 marks)

- **7** This question is about light.
 - (a) White light includes all the colours in the visible spectrum.

A beam of white light is the only light that shines on a book.

The book appears green.

A red filter is placed between the source of white light and the book.

What colour does the book appear now?

(1)

- A black
- **B** blue
- C green
- D red
- (b) Figure 8 shows a shiny metal ball.



(Source: © frerd/Shutterstock)

Figure 8

A clear image of a building can be seen on the surface of the ball.

- (i) This clear image is an example of
 - A diffuse reflection
 - **B** diffuse refraction
 - **C** specular reflection
 - **D** specular refraction

(1)



(ii)	Explain why the surface of the metal ball gives a clear image
	You may draw diagrams to help with your answer.

(2)

(c) Figure 9a and Figure 9b show rays of light before and after passing through different lenses.




Figure 9a




Figure 9b

(i) State **one** similarity and **one** difference in the way the rays of light pass through the lenses.

(2)

similarity.....

difference



(ii) Figure 10 shows two glass lenses P and Q.





Figure 10

State how the power of P is different from the power of Q.

(1)

*(d) Figure 11a shows refraction of light at a boundary between glass and air. Figure 11b shows total internal reflection of light at a boundary between glass and air.




glass

Figure 11a

Figure 11b

Use Figure 11a and Figure 11b to explain refraction and total internal reflection.

You may add to Figure 11a and Figure 11b to help with your answer.



8 (a) Figure 12 shows part of a wave.




Figure 12

Use data from Figure 12 to calculate the wavelength of the wave.

(2)

(b) (i) Figure 13 shows a student sitting on the shore of a lake watching ripples on the surface of the water moving past a toy boat.



Figure 13

The student has a stopwatch.

Describe how the student could determine the frequency of the ripples on the lake.

(3)

(ii) The speed of a water wave is 1.5 m/s.

The frequency of the wave is 0.70 Hz.

Calculate the wavelength of this wave.

Use the equation

$$v = f \times \lambda$$

(2)

wavelength = m



(iii) Water waves	are transverse	waves
-------------------	----------------	-------

Describe the difference between transverse waves and longitudinal waves.

(2)

(c) Sound waves travel at 330 m/s in air.

A student sees a flash of lightning.

The student hears the sound of thunder 4.0s later.

Calculate the distance from the student to the flash of lightning.

Use the equation

$$x = v \times t$$

(2)

distance = m

(Total for Question 8 = 11 marks)

9	This q	uestic	on is about radioactivity.	
	(a) Al	pha (d	α), beta (β) and gamma (γ) are three types of radioactive emissions.	
	W	hich s	statement describes all of these radioactive emissions?	(1)
	X	Α	ionising and emitted by stable nuclei	
	X	В	ionising and emitted by unstable nuclei	
	X	C	neutral and emitted by stable nuclei	
	X	D	neutral and emitted by unstable nuclei	
	(b) Flu	uorine	e-19 is a stable isotope of the element fluorine.	
	Th	ie elei	ment fluorine also has several radioactive isotopes.	
			e one similarity and one difference between the numbers of particles in	
	or	ne nuo	cleus of fluorine-19 and one nucleus of a radioactive isotope of fluorine.	(2)
sim	ilarity			
diff	erence	<u></u>		



(2)

(c) Figure 14 shows a Geiger-Muller (G-M) tube attached to a counter. The G-M tube is used to measure the activity of a source of beta (β) radiation. There is an aluminium sheet between the beta source and the G-M tube. The counter is switched on and after 1 minute shows a count of 268.




Figure 14

(i) The aluminium sheet is taken away.The counter is reset to zero and then switched on again.A new count is taken for 1 minute.

Explain why the new count is greater than 268.



The counter is reset to zero and switched on again. A new count is taken for 1 minute.	
Give a reason why there would now be a reading on the counter.	(1)
(iii) State the SI unit for the activity of a radioactive source.	(1)
Exposing people to radioactive sources can be dangerous.	
Describe the dangers of exposure to radioactive sources and what can be done to protect hospital staff when they are working with radioactive sources.	(6)
(Total for Question 9 = 13 ma	rke)





0 The Big Bang theory is one theory for the origin of the Universe.	
The Big Bang theory suggests:	
 the Universe had a beginning 	
the Universe is still expanding.	
(a) Which of these provides evidence that the Universe had a beginning?	(4)
■ A the discovery of other galaxies	(1)
■ B the discovery of the moons of Jupiter	
☑ the discovery of planets orbiting distant stars	
D the discovery of cosmic microwave background (CMB) radiation	
(b) Evidence that the Universe is still expanding comes from observations of lig from distant galaxies.	ht
Describe how light from distant galaxies shows that the Universe is still expand	anding.
(c) The Steady State theory is also a theory about the origin of the Universe.	
Give one similarity and one difference when comparing the Big Bang theory	y with
the Steady State theory.	(2)
	(2)
similarity	
difference	



(d) Observations of the expanding Universe have shown that the further away a galaxy is from the Earth, the faster the galaxy is moving away from the Earth.

Figure 15 shows how the velocity of galaxies is related to their distance from the Earth.




Figure 15

Mpc is a unit of distance used for large distances in space.

(i) Use Figure 15 to estimate the velocity of a galaxy that is 15 Mpc away from the Earth.

(1)

(ii) Calculate the gradient of the line shown in Figure 15.

State the unit.

(3)

(iii) The gradient of the line in Figure 15 can be used to estimate the Explain why the gradient of the line in Figure 15 can only provide of the age of the Universe.	_
of the age of the offiverse.	(2)
(Total for Questi	on 10 = 11 marks)
TOTAL FOR PA	PER = 100 MARKS

Equations

(final velocity)² – (initial velocity)² = $2 \times \text{acceleration} \times \text{distance}$

$$v^2 - u^2 = 2 \times a \times x$$

energy transferred = current \times potential difference \times time

$$E = I \times V \times t$$

potential difference across primary coil \times current in primary coil = potential difference across secondary coil \times current in secondary coil

$$V_p \times I_p = V_s \times I_s$$

change in thermal energy = mass \times specific heat capacity \times change in temperature

$$\Delta Q = m \times c \times \Delta \theta$$

thermal energy for a change of state = mass \times specific latent heat

$$Q = m \times L$$

to calculate pressure or volume for gases of fixed mass at constant temperature

$$P_1 V_1 = P_2 V_2$$

energy transferred in stretching = $0.5 \times \text{spring constant} \times (\text{extension})^2$

$$E = \frac{1}{2} \times k \times x^2$$

