

GCE

Physics A

H156/02: Depth in physics

Advanced Subsidiary GCE

Mark Scheme for November 2020

PMT

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2020

Here are the subject specific instructions for this question paper.

CATEGORISATION OF MARKS

The marking schemes categorise marks on the MACB scheme.

- **M** marks These are <u>method</u> marks upon which **A**-marks (accuracy marks) later depend. For an **M**-mark to be scored, the point to which it refers must be seen in the candidate's answers. If a candidate fails to score a particular **M**-mark, then none of the dependent **A**-marks can be scored.
- A marks These are accuracy or <u>answer</u> marks, which either depend on an **M**-mark, or allow a **C**-mark to be scored.
- **C** marks These are <u>compensatory</u> method marks which can be scored even if the points to which they refer are not written down by the candidate, providing subsequent working gives evidence that they must have known it. For example, if an equation carries a **C**-mark and the candidate does not write down the actual equation but does correct working which shows the candidate knew the equation, then the **C**-mark is given.
- **B** marks These are awarded as <u>independent</u> marks, which do not depend on other marks. For a **B**-mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.

SIGNIFICANT FIGURES

If the data given in a question is to 2 sf, then allow an answer to 2 or <u>more</u> significant figures. If an answer is given to fewer than 2 sf, then penalise once only in the <u>entire</u> paper. Any exception to this rule will be mentioned in the Guidance.

Annota	Annotations						
	Annotation	Meaning					
	Correct response	Used to indicate the point at which a mark has been awarded (one tick per mark awarded).					
×	Incorrect response	Used to indicate an incorrect answer or a point where a mark is lost.					
AE	Arithmetic error	Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.					
BOD	Benefit of doubt given	Used to indicate a mark awarded where the candidate provides an answer that is not totally satisfactory, but the examiner feels that sufficient work has been done.					
BP	Blank page	Use BP on additional page(s) to show that there is no additional work provided by the candidates.					
CON	Contradiction	No mark can be awarded if the candidate contradicts himself or herself in the same response.					
ECF	Error carried forward	Used in <u>numerical answers only</u> , unless specified otherwise in the mark scheme. Answers to later sections of numerical questions may be awarded up to full credit provided they are consistent with earlier incorrect answers. Within a question, ECF can be given for AE, TE and POT errors but not for XP.					
L1	Level 1	L1 is used to show 2 marks awarded and L1 [^] is used to show 1 mark awarded.					
L2	Level 2	L2 is used to show 4 marks awarded and L2 [^] is used to show 3 marks awarded.					
L3	Level 3	L3 is used to show 6 marks awarded and L3 [^] is used to show 5 marks awarded.					
РОТ	Power of 10 error	This is usually linked to conversion of SI prefixes. Do not allow the mark where the error occurs. Then follow through the working/calculation giving ECF for subsequent marks if there are no further errors.					
SEEN	Seen	To indicate working/text has been seen by the examiner.					
SF	Error in number of significant figures	Where more SFs are given than is justified by the question, do not penalise. Fewer significant figures than necessary will be considered within the mark scheme. Penalised only once in the paper.					
TE	Transcription error	This error is when there is incorrect transcription of the correct data from the question, graphical read-off, formulæ booklet or a previous answer. Do not allow the relevant mark and then follow through the working giving ECF for subsequent marks.					
ХР	Wrong physics or equation	Used in <u>numerical answers only</u> , unless otherwise specified in the mark scheme. Use of an incorrect equation is wrong physics even if it happens to lead to the correct answer.					
•	Omission	Used to indicate where more is needed for a mark to be awarded (what is written is not wrong but not enough).					

H156/02

PMT

Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

Annotation	Meaning
/	alternative and acceptable answers for the same marking point
Reject	Answers which are not worthy of credit
Not	Answers which are not worthy of credit
Ignore	Statements which are irrelevant
Allow	Answers that can be accepted
()	Words which are not essential to gain credit
	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument

Q	uesti	on	Answer	Marks	Guidance
1	(a)	(i)	Vibrations or oscillations parallel to direction of travel of the wave / direction of energy transfer	B1	
		(ii)	Amplitude of 2 cm (in each direction) Sinusoidal shape (by eye) with period of 4 cm – at least	B1 B1	Check peak, equilibrium and trough positions
	(1)	(1)	two waves	54	
	(b)	(i)	Microwave: 2 cm	B1	
			X-ray 200 pm	B1	
		(ii)	Any two from:	B1 x 2	
			May be reflected / refracted / diffracted / interference		
			May be polarised		
			Travel in a vacuum (at a constant speed / $3 \times 10^8 \text{m s}^{-1}$)		Allow speed of light
			Oscillation of electric and magnetic fields.		
			Total	7	

Q	Question		Answer	Marks	Guidance
2	(a)	(i)	$(f = v/\lambda) = 3.00 \times 10^8 \div 4.69 \times 10^{-7} (= 6.40 \times 10^{14} \text{ Hz})$	B1	6.397 x 10 ¹⁴ Hz
		(ii)	$1.96 \times 10^8 (\text{ms}^{-1})$	B1	
			3.07 × 10 ⁻⁷ (m)	B1	Allow 3.06×10^{-7} (m) (uses (i)) Not ECF for incorrect speed
	(b)		$p = 30^{\circ}$	B1	
		2.	$\sin q = 0.5 \times 1.53 \text{ or } 0.765$	C1	
			<i>q</i> = 50°	A1	Allow 49.9° Note 19 ° does not score
		(ii)	<i>p</i> always equals <i>i</i> or <i>p</i> increases with <i>i</i> / when $i = 60^\circ$, $p = 60^\circ$	B1	
			Any three from:	B1 x3	
			as <i>i</i> increases, <i>q</i> increases (until <i>i</i> equals the critical angle)		
			when $i = critical$ angle, $q = 90^{\circ}$		
			critical angle = 41°		
			when <i>i</i> is greater than critical angle, <u>total internal reflection</u> occurs		
			when $i = 60^{\circ}$, there is no angle q or no refracted ray		Not $q = 0$
	(c)		Straight line to centre of block and reflects along original ray P	B1	
			Straight line to centre of block and refracts with angle q less than 49.9° but greater than 30°	B1	
			Total	12	

Q	uesti	on	Answer	Marks	Guidance
3	(a)	(i)	points on the line read to the nearest half square	B1	Allow Δy for $y_2 - y_1$ and Δx for $x_2 - x_1$
			size of triangle is greater than half the length of the drawn line and Δy / Δx	B1	$\Delta x \ge 0.1625$
		(ii)	$\left(\frac{9.81}{0.12}\right) = 81.75$	C1	Allow ECF from (a)(i)
			82 N m ⁻¹ given to 2 or 3 significant figures	A1	Allow 81.8 N m ⁻¹ Note POT must be correct for given unit Allow kg s ⁻²
	(b)	(i)	steepest or shallowest line <u>that passes through all the</u> error bars	B1	
		(ii)	gradient determined: 0.10 mkg ⁻¹ or 0.13 mkg ⁻¹	B1	Allow ECF from (b)(i)
		(iii)	Δgradient (0.13 - 0.12 or 0.12 - 0.10)	C1	Allow ECF from (b)(i) and (ii)
			$\frac{\Delta \text{gradient}}{\text{gradient}} \times 100 = 8.3\% \text{ or } 17\%$	A1	Not 10% without justification
			OR	C1	
			Δk (82 – 75 or 98-82)		
			$\frac{\Delta k}{k} \times 100 = 8.5\%$ or 20%	A1	
			Total	8	

PMT

Question	Answer	Marks	Guidance
4	 Level 3 (5–6 marks) Detailed procedure including labelled diagram and measurements to be taken and detailed analysis There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated. Level 2 (3–4 marks) A diagram, some procedure, some measurements and some analysis or detailed analysis and limited procedure with limited diagram or or detailed procedure including diagram and measurements to be taken and limited analysis There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence. Level 1 (1–2 marks) Limited analysis There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant. 0 marks No response or no response worthy of credit.	B1 x 6	Indicative scientific points may include: Diagram and procedure • labelled diagram • horizontal surface supported • description of procedure • method to release ball • method to identify position ball hits the ground • repeats experiment for each v • method to prevent ball rolling on floor in laboratory Measurements • measuring instruments to determine v • measurements to determine $v e.g. mgh$ conversion or one light gate with diameter of ball measured or two light gates with distance between light gates measured • use of ruler to measure R . Analysis • equation to determine v • appropriate graph, e.g. plot R against v or plot R^2 against v^2 • Expect straight line passing through origin • $Q = g x$ gradient ² or $Q = g x$ gradient
	Total	6	

Q	uestion	Answer	Marks	Guidance
5	(a)	20 (m s ⁻¹)	B1	
	(b)	(After 0.75 s) gradient decreases with time	M1	
		Indicating velocity is decreasing/ deceleration	A1	
	(c)	Tangent drawn at $t = 1.75$ s (judge by eye)	B1	
		Gradient in the range 11.0 (m s ⁻¹) to 13.0 (m s ⁻¹)	B1	
	(d)	∆time = 1.75 – 0.75 OR 3.25 - 0.75	C1	Allow use of (c) and (a)
		20.12		Allow $a = 8.0 \text{ m s}^{-2}$ for $v^2 = u^2 + 2as$ or $s = ut + \frac{1}{2}at^2$ methods
		Using (c): $F = 950 \times \frac{20-12}{1.75-0.75}$ or	C1	Not ECF for incorrect time
		using graph: $F = 950 \times \frac{20-0}{3.25-0.75}$ or $F = \frac{950\times 20}{3.25-0.75}$		
		7600 (N)	A1	Ignore sign
	(e)	Maximum of two from:	B1 x 3	
		(thinking) time is the same		
		(braking) time is halved / 1.25 s		
		total time is 2 s		
		AND		
		maximum of two from:		
		(thinking) distance / displacement travelled (before braking) halved / 7.5 m		
		(braking) distance / displacement quarters / 6.25 m		
		total distance / displacement = 13.75 m		
		Total	11	

Q	uestic	on	Answer	Marks	Guidance
6	(a)		$\pi \times \frac{(32 \times 10^{-3})^2}{4} \times 100 \times 10^{-3} \text{ or } 8.04 \times 10^{-5}$	C1	Ignore POT
			$\frac{7.0}{9.81}$ or 0.714	C1	
			8900 (kg m ⁻³)	A1	8881 2200 scores two marks
	(b)	(i)	4.4 – 4.6 (N)	B1	
		(ii)	Weight of cylinder 3.5 cm vertically (judge by eye)	M1	
			Correct closed triangle drawn including T_A	M1	
			Correct directions indicated for weight and T_A and $T_A = 6.4 \pm 0.2$ (N)	A1	
		(iii)	39 ± 1°	A1	Allow ECF from (b)(ii) for trigonometry methods
	(c)		<i>F</i> x 100 or 7.0 x 16	C1	Ignore POT
			$F = \frac{7.0 x 16}{100} = 1.1 (\text{N})$	A1	1.12 Not 1.067
			Total	10	

PMT

Q	uesti	ion	Answer	Marks	Guidance
7	(a)	(i)	$R = \frac{150}{1.5^2}$	C1	Allow $V = \frac{150}{1.5} = 100 \text{ Vand } R = \frac{100}{1.5}$
			67 Ω	A1	
		(ii)	Q = 1.5 x 5.0 x 60 x 60 or 27000	C1	Note use of 150 (W) does not score 1.7 x 10 ²⁵
			$N = \frac{1.5 \times 5.0 \times 60 \times 60}{1.6 \times 10^{-19}} = 1.7 \times 10^{23}$	A1	1.68 x 10 ²³ 4.7 x 10 ¹⁹ scores one mark Not 1.7 x 10 ²⁵ (uses 150 W)
		(iii)	$v = \frac{1.5}{7.9 \times 10^{28} \times 4.1 \times 10^{-9} \times 1.6 \times 10^{-19}}$	C1	
			0.029 (m s ⁻¹)	A1	
	(b)		150 (x 10 ⁻³) x 5 x 16	C1	Not time in minutes or seconds
			12 (p)	A1	Allow ECF for POT on power
	(c)		Silicon will have a smaller number density, ORA	B1	Allow semiconductor for silicon; metal for nichrome
			Silicon will have a larger resistivity, ORA	B1	
			Total	10	

Question	Answer		Guidance
8	Level 3 (5–6 marks) Description and explanation of pattern changes and quantitatively explains link between de Broglie wavelength and potential difference. There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.	B1 x 6	 Indicative scientific points may include: Description of pattern changes Rings become closer (not just smaller) Rings become brighter Qualititative explanation of pattern changes in terms
	Level 2 (3–4 marks) Clear description of how pattern changes and explanation of pattern changes and qualitatively explains link between de Broglie wavelength and potential difference or	on en Electrons gain	of de Broglie wavelength and potential difference
	limited description of how pattern changes <u>and</u> quantitatively explains link between de Broglie wavelength and potential difference.		 Electrons have a greater momentum Implies smaller wavelength Smaller wavelength means less diffraction
	There is a line of reasoning presented with some structure. The information presented is in the most-part relevant and supported by some evidence.		 Shorter wavelength gives shorter path differences between areas of constructive and destructive interference Quantitative explanation of pattern changes in terms of de Broglie wavelength and potential difference eV = ¹/₂mv² p = mv
	Level 1 (1–2 marks) Limited description of how pattern changes and limited attempts to explain qualitatively the link between de Broglie wavelength and potential difference or		
	qualitatively explains link between de Broglie wavelength and potential difference.		
	The information is basic and communicated in an unstructured way. The information is supported by limited evidence and the relationship to the evidence may not be clear.		• $v^2 \alpha \text{ V or } p^2 \alpha \text{ V}$ • $\lambda = \frac{h}{p} \text{ or } \lambda \alpha \frac{1}{v}$ • $\lambda = \frac{h}{\sqrt{2meV}} \text{ or } \lambda \propto \frac{1}{\sqrt{v}}$
	0 marks No response or no response worthy of credit.		

OCR (Oxford Cambridge and RSA Examinations) The Triangle Building Shaftesbury Road Cambridge CB2 8EA

OCR Customer Contact Centre

Education and Learning Telephone: 01223 553998 Facsimile: 01223 552627 Email: <u>general.qualifications@ocr.org.uk</u>

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

